

**MOSQUITOES OF KUDAT: SPECIES COMPOSITION
AND THEIR MEDICAL IMPORTANCE
(DIPTERA: CULICIDAE)**

Ng S.H.¹, Homathevi R.² and Chua T.H.¹

¹Department of Pathobiology of Medical Diagnosis, Faculty of Medicine and Health Science, University Malaysia Sabah, Jalan UMS, 88400, Kota Kinabalu, Sabah.

²Institute Tropical Biology and Conservation, Universiti Malaysia Sabah, Jalan UMS, 88400, Kota Kinabalu, Sabah.

Corresponding author: thchua@ums.edu.my

ABSTRACT

A survey of mosquito fauna was done to determine the mosquito population and species composition in Kudat. Outdoor human landing catches (HLC) were done from April to June 2015. A total of 4,350 mosquitoes, representing eight genera and 37 species were collected. These include *Aedes* (*Stegomyia*), *Aedimorphus*, *Anopheles*, *Armigeres*, *Culex*, *Mansonia*, *Phagomyia* and *Uranotaenia*. Medical important species are specially noted and discussed. *Armigeres* (52.7%) was the most abundant, followed by *Culex* (31.9%), *Aedes* (9.4%), and *Anopheles* (4.3%). *Armigeres kesseli* and *Ar. subabaltus* were the two most common species. *Armigeres annulipalpi* and *Phagomyia prominens* are new records of Sabah.

Keywords: Mosquito species composition, new records, medical importance, Kudat

ABSTRAK

Kaji selidik berkaitan dengan fauna nyamuk telah dijalankan untuk mengenalpasti populasi nyamuk serta komposisi spesis yang terdapat di Kudat. Kaedah tangkapan berumpan manusia (“human landing catch”) telah dipakai dari bulan April 2015 sehingga Jun 2015. Sejumlah 4,350 nyamuk terdiri daripada lapan genera dan 37 spesis telah ditangkap. Antaranya ialah *Aedes* (*Stegomyia*), *Aedimorphus*, *Anopheles*, *Armigeres*, *Culex*, *Mansonia*, *Phagomyia* dan *Uranotaenia*. Spesis yang mempunyai kepentingan perubatan telah dibincang. Genus yang paling banyak termasuk *Armigeres* (52.7%), seterusnya *Culex* (31.9%), *Aedes* (9.4%), and *Anopheles* (4.3%). Spesis yang paling umum ialah *Armigeres kesseli* dan *Ar. subabaltus*. *Armigeres annulipalpi* and *Phagomyia prominens* ialah spesis rekod baru di Sabah.

Kata kunci: Spesies komposisi nyamuk, rekod baru, kepentingan perubatan, Kudat

INTRODUCTION

The importance of mosquitoes was highlighted when it was discovered as vector of filariasis in 1878 (Chernin, 1983), malaria in 1897 (Cox, 2010) and Japanese encephalitis in 1938 (Misra & Kalita, 2010). Mosquitoes are so important medically that it was considered as “the world deadliest animal”.

As one of the twelve mega-biodiversity hotspots of the world, Malaysia is a tropical country with a high number of mosquito species (Foley *et al.*, 2007). This environment also promotes the establishment of several endemic mosquito-borne diseases, including dengue, malaria, chikungunya, filariasis and Japanese encephalitis.

Many researchers have focused their researches in intervention of disease vector and overlooked fundamental studies such as vector biodiversity (Ferguson *et al.*, 2010). They seek more obvious and immediate potential benefits, but information on the ecology and bionomics of mosquitoes is essential to sustaining long term success. In fact such knowledge is a prerequisite to eradication or elimination of disease carrying mosquitoes and hence the disease themselves (Ferguson *et al.*, 2010). Knowledge of species composition is vital to provide database of mosquito diversity, which are medically important, so that limited resources could be applied effectively in their control. Thus, this study was conducted to find out the mosquito species composition.

MATERIALS AND METHOD

The study site was selected in Kudat District (6.71-7.26N, 116.75-117.22E), where the highest malaria cases of *Plasmodium knowlesi* in Malaysia were reported in 2011 (William *et al.*, 2013). Only five villages: Paradason, Marabahai, Tuboh, Nangka, Nangaran were included, with a total population of 640.

Mosquitoes were collected with outdoor human landing catch (HLC) as only anthropophilic species transmit pathogen to human. A total of 24 nights HLC from 1800-2400 hour were carried out in two weeks intervals from April 2015 – June 2015. For each night there were three teams of two collectors per team, stationed in three different sites randomly. Landing mosquitoes were collected with 2x2.5cm transparent plastic vials.

Collected mosquitoes were preserved and identified with published keys (Rattanaarithikul *et al.*, 2005; 2006a; 2006b). Literature search was carried out to identify all medical important species in collection.

The study was approved by the Medical Research & Ethic Committee of Ministry of Health Malaysia (NMRR-12-786-13048). All mosquito collectors signed informed consent forms and were provided with antimalarial prophylaxis.

RESULT

A total of 4,270 mosquitoes belonging to 36 species of 8 genera were collected (Table 1). The most abundant species was *Armigeres kesseli* (43.91%), followed by *Culex sitiens* (14.36%), *Aedes albopictus* (8.99%), *Ar. subabaltus* (7.56%), *Cx. vishnui* (3.19%), *Aedimorphus vexans* (2.69%), *Anopheles balabacensis* (2.58%), and *An. donaldi* (1.05%). The remaining species contributed less than 1%. Another 12.11% of mosquitoes belonging mostly to *Cx. vishnui* group were unidentifiable due to poor state of their morphological characteristics.

Among 36 species collected, 18 disease vectors were identified (Table 2). Human malaria is carried by anophelines: *An. balabacensis*, *An. donaldi*, and *An. maculatus*. Dengue, Chikungunya and Japanese encephalitis are carried by *Ae. aegypti* and *Ae. albopictus*. Japanese encephalitic is also transmitted by *Aedeomorphus vexans*, *Ar. subabaltus*, *Cx. bitaeniorhyncus*, *Cx. fuscocephala*, *Cx. quinquefasciatus*, *Cx. gelidus*, *Cx. sitiens*, *Cx. tritaeniorhyncus*, *Cx. vishnui*, *Cx. pseudovishnui*, and *Cx. whitmorei*. The primary vectors of filariasis are *Mansonia uniformis* and *Man. indiana* while secondary vectors are *Cx. quinquefasciatus*, *An. balabacensis*, *An. donaldi*, *Ar. subabaltus*, and *Coquillettidia crassipes*.

DISCUSSION

Armigeres subabaltus and *Ar. kesseli* were found to be dominant most likely due to the abundance of breeding sites. Many *Armigeres* larvae were found breeding in many coconut shells during site investigation. The coconut shell can also be

mutual breeding sites for *Armigeres jugraensis*, *Ar. malayi* and *Ar. moultoni* (Toma *et al.*, 2012). Besides, the improper management of sewage system, such as the uncovered septic tank provides ideal breeding sites to *Armigeres* spp (Rajavel, 1992). This putrefied water found in sewage and agricultural plantation could be helped the breeding of *Culex sitiens*, as it was also found in southern Thailand (Prummongkol *et al.*, 2012).

Larvae of *Aedes albopictus* were abundant in water drums and containers. Some third instar larvae were also found breeding in a tree holes of forested area. These explains the plasticity of *Ae. albopictus* in breeding site selection (Gratz, 2004).

Another finding is the host seeking behaviour of *Ae. albopictus*, a diurnal species (Sullivan *et al.*, 1971), however, our result suggested that *Ae. albopictus* is also active at night when host approaches (Dieng *et al.*, 2015). We also collected three adults of *Ae. aegypti*. These two species are very competent in arboviruses transmission (Gratz, 2004) and are the most important dengue vectors in Malaysia. Thus their presence could be a risk factor to local people.

Anopheles balabacensis is primary human malaria vector in northern Borneo (Reid, 1968). It is the predominant anopheline in Kudat where the other species are represented in a small percentage (Wong *et al.*, 2015). *An. donaldi* and *An. maculatus* are the malaria vector in Sandakan (Goh *et al.*, 2013) and Peninsula Malaysia (Reid, 1968). *Armigeres subabaltus* has been recorded positive with *Brugia pahangi* filarial worms in Peninsula Malaysia (Muslim *et al.*, 2013), but there is no evidence to show that it is a disease vector in Borneo island. The vectorial potential of *Armigeres* mosquitoes should be determined in future study.

Finally, *Armigeres annulipalpi* and *Phagomyia prominens* were recorded for the first time in Sabah. These two species are distributed in Oriental area (VectorMap, 2014) such as China, India, Vietnam, Cambodia, Nepal, Thailand and Indonesia. The vector status of these two species in Sabah is currently unknown.

ACKNOWLEDGEMENT

This study was carried out as a minor assignment by entomology component of the Monkeybar Malaria Project funded by London School of Hygiene & Tropical Medicine. The effort of local villagers involved in mosquito catches is very much appreciated.

Table 1 Species composition of mosquitoes

Species	Apr	May	Jun	%
<i>Aedimorphus vexans</i> (Meigen)	-	-	115	2.69
<i>Anopheles balabacensis</i> Baisas	51	19	40	2.58
<i>Anopheles donaldi</i> Reid	18	10	17	1.05
<i>Anopheles maculatus</i> Theobald	11	-	-	0.26
<i>Anopheles umbrosus</i> gp.	8	1	-	0.21
<i>Armigeres annulipalpis</i> (Theobald)	1	-	1	0.05
<i>Armigeres jugaensis</i> (Leicester)	-	15	6	0.49
<i>Armigeres kesseli</i> Ramanlingam	645	565	665	43.91
<i>Armigeres kuchingensis</i> Edwards	-	2	-	0.05
<i>Armigeres malayi</i> (Theobald)	-	2	-	0.05
<i>Armigeres moultoni</i> Edwards	1	-	-	0.02
<i>Armigeres subabaltus</i> (Coquillett)	99	94	130	7.56
<i>Coquillettidia crassipes</i> (van der Wulp)	-	1	-	0.02
<i>Culex alis</i> Theobald	5	-	-	0.12
<i>Culex bitaeniorhynchus</i> Giles	-	-	3	0.07
<i>Culex gelidus</i> Theobald	1	5	13	0.44
<i>Culex pseudovishnui</i> Colless	-	4	6	0.23
<i>Culex quinquefasciatus</i> Say	7	5	7	0.44
<i>Culex sitiens</i> Wiedemann	18	445	150	14.36
Total	865	1168	1153	

Species	Apr	May	Jun	%
<i>Culex tritaeniarhycus</i> Giles	5	3	8	0.37
<i>Culex vishnui</i> Theobald	9	37	90	3.19
<i>Culex whitei</i> Barraud	-	1	-	0.01
<i>Culex whitmorei</i> (Giles)	-	1	-	0.01
<i>Cx. sp1</i>	121	112	284	12.11
<i>Mansonia indiana</i> Edwards	-	1	-	0.01
<i>Mansonia uniformis</i> (Theobald)	1	1	8	0.23
<i>Mansonia sp1</i>	-	1	-	0.05
<i>Paraaedes sp1</i>	-	1	-	0.05
<i>Phagomyia prominens</i> (Barraud)	-	-	1	0.05
<i>Aedes aegypti</i> (Linnaeus)	-	-	3	0.07
<i>Aedes albopictus</i> (Skuse)	5	217	162	8.99
<i>Aedes malayensis</i> (Colless)	-	1	-	0.05
<i>Aedes pseudalbopicta</i> Borel	2	2	2	0.14
<i>Aedes patriciae</i> (Mattingly)	-	2	-	0.05
<i>Aedes w-albus</i> Theobald	-	-	1	0.01
<i>Aedes sp1</i>	-	1	-	0.01
<i>Uranotaenia lateralis</i> Ludlow	-	1	-	0.01
Total	143	382	559	
Grand total				4,270/100%

Table 2 Medical important mosquito species

Species	Medical important	References
<i>Aedeomorphus vexans</i>	Japanese Encephalitis	Sucharit <i>et al.</i> , 1989
<i>Aedes aegyti</i> <i>Aedes albopictus</i>	Chikungunya	Noridah <i>et al.</i> , 2007; Pulmanusahakul <i>et al.</i> , 2011
	Dengue	Singh & Paul, 1969; Gratz, 2004; Lee & Rohani, 2005
	JE	Misra & Kalita, 2010; Vythilingam <i>et al.</i> , 1995
<i>Anopheles balabacensis</i>	Malaria	Reid, 1968; Hii <i>et al.</i> , 1985; Wong <i>et al.</i> , 2015
	Filariasis	Hii <i>et al.</i> , 1985
<i>Anopheles donaldi</i>	Malaria	Vythilingam <i>et al.</i> , 2005
	Filariasis	Reid, 1968; Vythilingam <i>et al.</i> , 1996
<i>Anopheles maculatus</i>	Malaria	Reid, 1968
<i>Armigeres subabaltus</i>	Filariasis	Aliota <i>et al.</i> , 2011; Muslim <i>et al.</i> , 2013
	JE	Rosen, 1986; Misra & Kalita, 2010
<i>Coquillettidia crassipes</i>	Filariasis	Chiang <i>et al.</i> , 1984
<i>Culex.bitaeniorhycus</i>	JE	Sucharit <i>et al.</i> , 1989; Vythilingam <i>et al.</i> , 1995
<i>Culex gelidus</i>	JE	Macdonald <i>et al.</i> , 1967; Vythilingam <i>et al.</i> , 1997
<i>Culex pseudovishnui</i>	JE	Sucharit <i>et al.</i> , 1989
<i>Culex quinquefasciatus</i>	JE	Rosen, 1986; Vythilingam <i>et al.</i> , 1997; Misra & Kalita, 2010
	Filariasis	Vythilingam <i>et al.</i> , 2005
<i>Culex sitiens</i>	JE	Vythilingam <i>et al.</i> , 1995; Vythilingam <i>et al.</i> , 2002
<i>Culex tritaeniorhycus</i>	JE	Macdonald <i>et al.</i> , 1967; Vythilingam <i>et al.</i> , 1995; 1997
<i>Culex vishnui</i>	JE	Sucharit <i>et al.</i> , 1989; Vythilingam <i>et al.</i> , 1995
<i>Culex whitmorei</i>	JE	Peiris <i>et al.</i> , 1993
<i>Mansonia indiana</i>	Filariasis	Wharton, 1962; Chiang, 1993
<i>Mansonia uniformis</i>	Filariasis	Wharton, 1962; Chiang <i>et al.</i> , 1984; Chiang, 1993

REFERENCES

- Aliota, M. T., Chen, C.-C., Dagoro, H., Fuchs, J. F., & Christensen, B. M. 2011. Filarial Worms Reduce Plasmodium Infectivity in Mosquitoes. *PLoS Neglected Tropical Diseases*, 5(2), 963.
- Chernin, E. 1983. Sir Patrick Manson's studies on the transmission and biology of filariasis. *Reviews of Infectious Diseases*, 5(1), 148–66.
- Chiang, G. L. 1993. Update on the bionomics of *Mansonia* vectors of brugian filariasis. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 24 Suppl 2, 69–75.
- Chiang, G. L., Cheong, W. H., Samarawickrema, W. A., Mak, J. W., & Kan, S. K. 1984. Filariasis in Bengkoka Peninsula, Sabah, Malaysia: vector studies in relation to the transmission of filariasis. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 15(2), 179–89.
- Cox, F. E. G. 2010. History of the discovery of the malaria parasites and their vectors. *Parasites & Vectors*, 3(5), 1–9.
- Dieng, H., Hui, O. S., Hassan, A. A., Abang, F., Ghani, I. A., Satho, T., Nolasco-Hipolito, C. 2015. Changes in the biting activity of a dengue vector relative to larval and adult nutritional histories: Implications for preventive measures. *Journal of Asia-Pacific Entomology*, 18(3), 507–513.
- Ferguson, H. M., Dornhaus, A., Beeche, A., Borgemeister, C., Gottlieb, M., Mulla, M. S., Killeen, G. F. 2010. Ecology: a prerequisite for malaria elimination and eradication. *PLoS Medicine*, 7(8).
- Foley, D. H., Rueda, L. M., & Wilkerson, R. C. 2007. Insight into global mosquito biogeography from country species records. *Journal of Medical Entomology*, 44(4), 554–67.

- Goh, X. T., Lim, Y. A. L., Vythilingam, I., Chew, C. H., Lee, P. C., Ngui, R., Chua, K. H. 2013. Increased detection of *Plasmodium knowlesi* in Sandakan division, Sabah as revealed by PlasmoNex™. *Malaria Journal*, 12(1), 264.
- Gratz, N. G. 2004. Critical review of the vector status of *Aedes albopictus*. *Medical and Veterinary Entomology*, 18(3), 215–227.
- Hii, J., Kan, S., Vun, Y. S., Chin, K. F., Lye, M. S., Make, J. W., & Cheong, W. H. 1985. *Anopheles flavirostris* incriminated as a vector of malaria and Bancroftian filariasis in Banggi Island, Sabah, Malaysia. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 79, 677–680.
- Lee, H. L., & Rohani, A. 2005. Transovarial transmission of dengue virus in *Aedes aegypti* and *Aedes albopictus* in relation to dengue outbreak in an urban area in Malaysia. *Dengue Bulletin*, 29, 106–111.
- Macdonald, W. W., Smith, C. E., Dawson, P. S., Ganapathipillai, A., & Mahadevan, S. 1967. *Arbovirus* infections in Sarawak: further observations on mosquitoes. *Journal of Medical Entomology*, 4(2), 146–57.
- Misra, U. K., & Kalita, J. 2010. Overview: Japanese encephalitis. *Progress in Neurobiology*, 91(2), 108–120.
- Muslim, A., Fong, M.-Y., Mahmud, R., Lau, Y.-L., & Sivanandam, S. 2013. *Armigeres subalbatus* incriminated as a vector of zoonotic *Brugia pahangi* filariasis in suburban Kuala Lumpur, Peninsular Malaysia. *Parasites & Vectors*, 6(1), 219.
- Noridah, O., Paranthaman, V., Nayar, S. K., Masliza, M., Ranjit, K., Norizah, I., Chua, K. B. 2007. Outbreak of chikungunya due to virus of Central/East African genotype in Malaysia. *Medical Journal of Malaysia*, 62(4), 323–328.

- Peiris, J. S. M., Amerasinghe, F. P., Arunagiri, C. K., Perera, L. P., Karunaratne, S. H. P. P., Ratnayake, C. B., Abeysinghe, M. R. N. 1993. Japanese encephalitis in Sri Lanka: comparison of vector and virus ecology in different agro-climatic areas. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 87(5), 541–548.
- Prummongkol, S., Panasoponkul, C., Apiwathnasorn, C., & Lek-Uthai, U. 2012. Biology of *Culex sitiens*, a predominant mosquito in Phang Nga, Thailand after a tsunami. *Journal of Insect Science (Online)*, 12(11),.
- Pulmanausahakul, R., Roytrakul, S., Auewarakul, P., & Smith, D. R. 2011. Chikungunya in Southeast Asia: understanding the emergence and finding solutions. *International Journal of Infectious Diseases*, 15(10), 671–676.
- Rajavel, A. R. 1992. Larval habitat of *Armigeres subalbatus* (COQ) and its characteristics in Pondicherry. *Southeast Asian J Trop Med Public Health*, 23(3), 470–473.
- Rattanarithikul, R., Harbach, R. E., Harrison, B. A., Panthusiri, P., Jones, J. W., Pelton, E., & Coleman, R. 2005. Illustrated Key to The Mosquito of Thailand II. Genera *Culex* and *Lutzia*. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 36 (supplément 2).
- Rattanarithikul, R., Harrison, B. A., Harbach, R. E., Panthusiri, P., & Coleman, R. E. 2006. Illustrated keys to the mosquitoes of Thailand-IV. *Anopheles*. *Southeast Asian J Trop Med Public Health*, 37, 1–128.
- Rattanarithikul, R., Harrison, B. A., Panthusiri, P., Peyton, E. L., & Coleman, R. E. 2006. Illustrated keys to the mosquitoes of Thailand: III. Genera *Aedeomyia*, *Ficalbia*, *Mimomyia*, *Hodgesia*, *Coquillettia*, *Mansonia*, and *Uranotaenia*. *Southeast Asian Journal of Tropical Medicine and Public Health*, 37(SUPPL. 1), 1–10.
- Reid, J. A. 1968. *Anopheles* of Malaya and Borneo. *Studies from the Institute for Medical Research Malaysia No. 31*.

- Rosen, L. 1986. The natural history of Japanese encephalitis virus. *Annu Rev Microbiol*, 40(52), 395–414.
- Singh, K. R., & Paul, S. D. 1969. Isolation of Dengue viruses in *Aedes albopictus* cell cultures. *Bulletin of the World Health Organization*, 40(6), 982–3.
- Sucharit, S., Surathin, K., & Shrestha, S. R. 1989. Vectors of Japanese encephalitis virus (JEV): species complexes of the vectors. *The Southeast Asian Journal of Tropical Medicine and Public Health*, 20(4), 611–21.
- Sullivan, M. F., Gould, D. J., & Maneechai, S. 1971. Observations on the host range and feeding preferences of *Aedes albopictus* (Skuse). *J Med Entomol*, 8(6), 713–716.
- Toma, T., Miyagi, I., Okazawa, T., Higa, Y., Wong, S. F., Leh, M. U., & Yong, H. Sen. 2012. *Redescriptions of Armigeres annulipalpis and Armigeres flavus (Diptera: Culicidae) from Sarawak, Malaysia*, 5–19.
- VectorMap. 2014. Mosquito species country list. Retrieved from <http://vectormap.si.edu/MosquitoCountryList.htm>
- Vythilingam, I., Chan, S. T., Shanmugratnam, C., Tanrang, H., & Chooi, K. H. 2005. The impact of development and malaria control activities on its vectors in the Kinabatangan area of Sabah , East Malaysia, 96, 24–30.
- Vythilingam, I., Lokman, H., ST, C., & JW, M. 1996. *Anopheles donaldi* incriminated as a vector of filariasis.
- Vythilingam, I., Oda, K. ., Chew, T. K. ., Mahadevan, S.; Vijayamalar, B. ., Morita, K. ., Tsuchie, H. ., & Igarashi, A. 1995. Isolation of Japanese encephalitis virus from mosquitoes collected in Sabak Bernam, Selangor, Malaysia in 1992. *Journal of the American Mosquito Control Association*, 11(1), 94.

- Vythilingam, I., Oda, K., Mahadevan, S., Abdullah, G., Thim, C. S., Hong, C. C., Igarashi, a. 1997. Abundance, parity, and Japanese encephalitis virus infection of mosquitoes (Diptera: Culicidae) in Sepang District, Malaysia. *Journal of Medical Entomology*, 34(3), 257–262.
- Vythilingam, I., Tan, C. H., & Nazni, W. A. 2005. Transmission potential of *Wuchereria bancrofti* by *Culex quinquefasciatus* in urban areas of Malaysia. *Tropical Biomedicine*, 22(1), 83–5.
- Vythilingam, I., Tan, S. B., & Krishnasamy, M. 2002. Short communication: Susceptibility of *Culex sitiens* to Japanese encephalitis virus in peninsular Malaysia. *Tropical Medicine & International Health*, 7(6), 539–540.
- Wharton, R. H. 1962. The biology of *Mansonia* mosquitoes in relation to the transmission of filariasis in Malaya. *Bulletin - Institute for Medical Research, Kuala Lumpur*, 11, 1–114.
- William, T., Rahman, H. A., Jelip, J., Ibrahim, M. Y., Menon, J., Grigg, M. J., Barber, B. E. 2013. Increasing Incidence of *Plasmodium knowlesi* Malaria following Control of *P. falciparum* and *P. vivax* Malaria in Sabah, Malaysia. *PLoS Neglected Tropical Diseases*, 7.
- Wong, M. L., Chua, T. H., Leong, C. S., Khaw, L. T., Fornace, K., Wan-Sulaiman, W.-Y., Vythilingam, I. 2015. Seasonal and Spatial Dynamics of the Primary Vector of *Plasmodium knowlesi* within a Major Transmission Focus in Sabah, Malaysia. *PLoS Neglected Tropical Diseases*, 9(10).